

ON 2-DIMENSIONAL NONASPHERICAL CELL-LIKE PEANO CONTINUA: A SIMPLIFIED APPROACH

KATSUYA EDA, UMED H. KARIMOV, AND DUŠAN REPOVŠ

ABSTRACT. We construct a functor $AC(-, -)$ from the category of path connected spaces X with a base point x to the category of simply connected spaces. The following are the main results of the paper: (i) If X is a Peano continuum then $AC(X, x)$ is a cell-like Peano continuum; (ii) If X is n -dimensional then $AC(X, x)$ is $(n + 1)$ -dimensional; and (iii) For a path connected space X , $\pi_1(X, x)$ is trivial if and only if $\pi_2(AC(X, x))$ is trivial. As a corollary, $AC(S^1, x)$ is a 2-dimensional nonaspherical cell-like Peano continuum.

1. INTRODUCTION

Peano continua are those continua (i.e. connected metric compacta) which are also locally connected. This classical topics of topology has received renewed attention in recent years, in particular the constructions of some surprising examples which provide answers to some very interesting problems and conjectures.

In our investigations of Peano continua we constructed in [6] a new functor from the category of all path connected spaces X with a base point x and continuous mappings to the category of all simply connected spaces, which we named the *Snake cone* and denoted by $SC(-, -)$. When X is a Peano continuum, $SC(X, x)$ is also a Peano continuum. We proved that in the case when $X = S^1$, the space $SC(S^1, x)$ is a noncontractible simply connected cell-like 2-dimensional Peano continuum.

In a subsequent paper [7] we showed that the space $SC(S^1, x)$ is even nonaspherical, i.e. $\pi_2(SC(S^1, x)) \neq 0$. Moreover, we showed that for every path connected space X , $\pi_1(X, x)$ is trivial if and only if $\pi_2(SC(X, x))$ is trivial (cf. [7] and [8]).

In the present paper we construct yet another new useful functor, called the *Alternating cone*, and we denote it by $AC(-, -)$. We show that while $AC(X, x)$ shares all the properties of the space $SC(X, x)$ listed above, their verification is much easier. We also show that there exists a Peano continuum (X, x) such that the spaces $AC(X, x)$ and $SC(X, x)$ are not homotopy equivalent. As an example how much easier the verifications are for the new functor, we consider singular homology $H_*(AC(X, x))$ (cf. Theorem 3.4).

2. PRELIMINARIES

We start by fixing some terminology and notations which will be used in the sequel. All undefined terms can be found in [13] or [14].

Date: February 19, 2013.

2010 Mathematics Subject Classification. Primary 54F15, 55N15; Secondary 54G20, 57M05.

Key words and phrases. Noncontractible compactum, weak homotopy equivalence, trivial shape, Peano continuum, Snake cone, Alternating cone, asphericity, cell-like space, Topologist sine curve.

For any two points A and B of the plane \mathbb{R}^2 , we shall denote the *linear segment* connecting these points by $[A, B]$. Next, by the *open* (resp. *half-open*) *interval* we shall mean the linear segment without its end points (resp. without one of its end points), and we shall denote it by (A, B) (resp. $[A, B)$ or $(A, B]$), as usual. The unit segment of the real line \mathbb{R}^1 , with the natural topology, will be denoted by \mathbb{I} . The point of the coordinate plane \mathbb{R}^2 with coordinates a and b will be denoted by $(a; b)$.

Consider the points $A = (0; 0)$, $B = (0; 1)$, $A_n = (1/n; 0)$, $B_n = (1/n; 1)$ and let $L = [A, B]$, $L_{2n-1} = [A_n, B_n]$, $L_{2n} = [B_n, A_{n+1}]$ be segments in the plane \mathbb{R}^2 , for $n \in \mathbb{N} = \{1, 2, 3, \dots\}$. The piecewise linear *Topologist sine curve* \mathcal{T} is a subspace of the plane \mathbb{R}^2 defined as the union of L_n and L . For any topological space X the cone $C(X)$ over X is defined as the quotient space $C(X) = (X \times \mathbb{I})/(X \times \{1\})$.

Suppose that $q_i : Z_i \rightarrow \mathbb{I}$, $i = 1, 2$, are mappings of topological spaces. Then the mapping $f : Z_1 \rightarrow Z_2$ is said to be *flat* with respect to the mappings q_1 and q_2 if for every two points $a, b \in Z_1$ with $q_1(a) = q_1(b)$, the equality $q_2(f(a)) = q_2(f(b))$ holds. Homotopy $H : Z_1 \times \mathbb{I} \rightarrow Z_2$ is said to be *flat* if for each $t \in \mathbb{I}$ the mapping $H(-, t) : Z_1 \rightarrow Z_2$ is a flat mapping with respect to the mappings q_1 and q_2 .

We recall the notion of the free σ -product of groups and a lemma from [4] (cf. the proof of [7, Theorem 3.1]). We only use it in the case when the index set is countable and we introduce a restricted version in the sequel. Let (X_n, x_n) be pointed spaces for $n \in \mathbb{N}$ such that $X_m \cap X_n = \emptyset$, for $m \neq n$. The underlying set of the pointed space $(\widetilde{\bigvee}_{n \in \mathbb{N}}(X_n, x_n), x^*)$ is the union of all X_n 's obtained by identifying all x_n 's to a point x^* and the topology is defined by specifying the neighborhood base as follows (cf. [1]):

- (1) If $x \in X_n \setminus \{x_n\}$, then the neighborhood base of x in $\widetilde{\bigvee}_{n \in \mathbb{N}}(X_n, x_n)$ is the one of X_n ; and
- (2) The point x^* has a neighborhood base, each element of which is of the form: $\bigcup_{n \geq m}(X_n, x_n) \cup \bigcup_{n < m} U_n$ where $m \in \mathbb{N}$ and each U_n is an open neighborhood of x_n in X_n for $n < m$.

Lemma 2.1. [4, Theorem A.1] *Let X_n be locally simply-connected and first countable at x_n for each n . Then*

$$\pi_1(\widetilde{\bigvee}_{n \in \mathbb{N}}(X_n, x_n), x^*) \cong \times_{n \in \mathbb{N}} \pi_1(X_n, x_n).$$

Let G be a group. A *commutator* is an element $[g_1, g_2] = g_1 g_2 g_1^{-1} g_2^{-1}$ where $g_1, g_2 \in G$. The normal subgroup G' of G which is generated by all commutators is called the *commutator subgroup* of G .

Recall that the *commutator length* $cl(g)$ of $g \in G'$ is defined as the minimal number of the commutators of G whose product is equal to g (cf. [5]). We note that $cl(e) = 0$ for the identity element e of the group G . If $\varphi : G \rightarrow H$ is a homomorphism of a group G to a group H , then for every $g \in G'$,


$$(1) \quad cl(\varphi(g)) \leq cl(g).$$

For every path connected space X , the fundamental group $\pi_1(X, x)$ is independent of the choice of the base point x and hence we can simply write $\pi_1(X)$. Finally, recall the isomorphism:

$$(2) \quad H_1(X) \cong \pi_1(X)/\pi'_1(X).$$

3. CONSTRUCTIONS AND MAIN RESULTS


The Snake cone functor $SC(-, -)$ is defined as follows. For every compact space X with the base point x , the space $SC(X, x)$ is the quotient space of the topological sum $(\mathcal{T} \times X) \sqcup \mathbb{I}^2$ via the identification of the points $(t, x) \in \mathcal{T} \times X$ with $t \in \mathcal{T} \setminus L \subset \mathbb{I}^2$ and the identification of each set $\{t\} \times X$ with the point t , for every $t \in L$ (cf. [6]).

FIGURE 1. $SC(S^1)$

Let us now define the Alternating cone functor $AC(-, -)$. Let (X, x) be a pointed space and let (X_n, x_n) be its copies. Connect x_n and x_{n+1} by an arc for each $n \in \mathbb{N}$ and add a new point so that a neighborhood of the point contains almost all X_n 's and almost all attached arcs. Let Y be this space.

Let $AC(X, x)$ be the quotient space of $Y \times \mathbb{I}$ in which the sets $\{0\} \times X_n$ are identified with $(0, x_n)$ for all odd n and the sets $\{1\} \times X_n$ are identified with $(1, x_n)$ for all even n .

Since x_n and the attached arcs converge to the new point in Y , $AC(X, x)$ can be illustrated as in Figure 2. The identified points will be denoted by $A_{2n} = (1/2n; 0)$ and $B_{2n-1} = (1/(2n-1); 1)$. Let $p : AC(X, x) \rightarrow \mathbb{I}^2$ be the natural projection and define $p(u) = (p_1(u); p_2(u))$. We let $C_n = (1/n; 1/2)$, for $n \in \mathbb{N}$ and $C = (0; 1/2)$.

FIGURE 2. $AC(S^1)$

Proposition 3.1. *Let X be a path connected space and $x, y \in X$. Then $SC(X, x)$ and $SC(X, y)$ are homotopy equivalent. Similarly, $AC(X, x)$ and $AC(X, y)$ are also homotopy equivalent.*

Proof. We shall only prove this for $SC(X, x)$ and $SC(X, y)$, since the proof for $AC(X, x)$ and $AC(X, y)$ is similar. Let X be a path connected space with $x \in X$, and X^* the space defined in Theorem 3.4. Also, let Y^* be the space obtained by replacing x with y . By a deformation on \mathbb{I}^2 , $SC(X, x)$ is homotopy equivalent to $SC(X^*, 0)$. Hence it suffices to establish a homotopy equivalence between $SC(X^*, 0)$ and $SC(Y^*, 0)$.

Let $f : \mathbb{I} \rightarrow X$ be a path such that $f(0) = x$ and $f(1) = y$. Define $\varphi : X^* \rightarrow Y^*$ and $\psi : Y^* \rightarrow X^*$ by:

$$\varphi(u) = \begin{cases} u & \text{for } u \in X \\ 2s & \text{for } s \in [0, 1/2] \\ f(2 - 2s) & \text{for } s \in [1/2, 1] \end{cases}$$

$$\psi(u) = \begin{cases} u & \text{for } u \in X \\ 2s & \text{for } s \in [0, 1/2] \\ f(2s - 1) & \text{for } s \in [1/2, 1]. \end{cases}$$

Then it is easy to see that φ and ψ induce a homotopy equivalence between X^* and Y^* which fixes 0. This homotopy equivalence then induces a homotopy equivalence between $SC(X^*, 0)$ and $SC(Y^*, 0)$. \square

We are mainly interested in the homotopy types of $SC(X, x)$ and $AC(X, x)$ for the class of path connected spaces X . Since by Proposition 3.1 the choice of the base point x does not affect the homotopy types of $SC(X, x)$ and $AC(X, x)$, we can simply write $SC(X)$ or $AC(X)$ when the choice of the base point does not matter.

Theorem 3.2. *If X is a continuum then $AC(X)$ is a cell-like space. If $\dim X = n$ then $\dim AC(X) = n + 1$. If X is a Peano continuum, then $AC(X)$ is also a Peano continuum. For any continuum X , the space $AC(X)$ is simply connected.*

Theorem 3.3. *If X is a noncontractible compact space, then $AC(X)$ is also a noncontractible compact space.*

Theorem 3.4. *Let X be a path connected space with $x \in X$ and X^* the quotient space of the disjoint sum of the unit interval \mathbb{I} and X , by identifying 1 of \mathbb{I} and x . Let (X_i, x_i) be copies of $(X^*, 0)$.*

It then follows that for any natural number n , we have the isomorphism:

$$H_{n+1}(AC(X)) \cong H_n(C(\tilde{\bigvee}_{i \in \mathbb{N}} (X_{2i-1}, x_{2i-1})) \vee C(\tilde{\bigvee}_{i \in \mathbb{N}} (X_{2i}, x_{2i}))),$$

where the attaching points of the two cones are the identification points x_{2i-1} and x_{2i} .

Theorem 3.5. *Let X be a path connected space. Then $\pi_1(X)$ is trivial if and only if $\pi_2(AC(X))$ is trivial.*

It is easy to see that if $\{a, b\}$ is a two-point space with the discrete topology then the space $AC(\{a, b\})$ is path connected, whereas the space $SC(\{a, b\})$ is not. The following theorem shows their difference from another perspective:

Theorem 3.6. *There exists a Peano continuum X such that $AC(X)$ is not homotopy equivalent to $SC(X)$.*

4. PROOFS OF THEOREMS 3.2 AND 3.3

In this section we shall prove Theorems 3.2 and 3.3. The argument is similar to the one which was used for proving analogous statements concerning $SC(X)$.

Proof of Theorem 3.2. Since X is a compactum, the space $AC(X)$ is the inverse limit of an inverse sequence of contractible compact metrizable spaces. Therefore $AC(X)$ is a compact metrizable space of trivial shape, i.e. cell-like.

If $\dim X = n$ and X is compact and metrizable then the dimension of its one-point compactification Y is equal to n and thus $\dim(Y \times \mathbb{I}) = n + 1$. Therefore $\dim AC(X) = n + 1$. If X is a Peano continuum then $AC(X)$ is compact, metrizable and locally path connected, hence also a Peano continuum.

Let us prove that $AC(X)$ is a simply connected space. Let $U = p_2^{-1}([0, 1])$ and $V = p_2^{-1}((0, 1])$ (cf. Figure 1). The sets U and V are homotopy equivalent to $p_2^{-1}(\{0\})$ and $p_2^{-1}(\{1\})$, respectively. Their intersection $U \cap V$ is homotopy equivalent to $p_2^{-1}(\{1/2\})$. Obviously, the mapping

$$\pi_1(p_2^{-1}(\{1/2\})) \rightarrow \pi_1(p_2^{-1}(\{0\})) * \pi_1(p_2^{-1}(\{1\}))$$

is surjective and the intersection $U \cap V$ is path connected. Therefore by the van Kampen theorem, the union $U \cup V = AC(X)$ is simply connected. \square

For a noncontractible space X , the noncontractibility of $AC(X)$ can be proved similarly as for $SC(X)$ (cf. [6, Theorem 1.2]). Hence we only indicate the difference and we state the necessary lemmas.

Let $M_y = p_2^{-1}(\{y\})$ and $AC_n(X) = p^{-1}([A_{2n}, B_{2n}] \cup [A_{2n+1}, B_{2n+1}])$ be the subspaces of $AC(X)$. The following lemma can be proved analogously as [6, Lemma 4.2].

Lemma 4.1. *Let $n \in \mathbb{N}$ and let $H : AC_n(X) \times \mathbb{I} \rightarrow AC(X)$ be a mapping such that for every $y \in \mathbb{I}$ and $t \in \mathbb{I}$, the closure of the set $p_2(H(M_y \cap AC_n(X), t))$ does not contain both points 0 and 1, and such that both mappings $H(-, 0)$ and $H(-, 1)$ are flat. Then there exists a flat homotopy from $H(-, 0)$ to $H(-, 1)$.*

For $s \in (0, 1)$ and $t \in \mathbb{I}$, we define a property $P(s, t)$ of the flat homotopy H as follows:

$$\begin{aligned} &H(M_s \cap AC_n(X), t) \subseteq p_2^{-1}((0, 1)) \text{ and the restriction of } H(-, t) \text{ to} \\ &M_s \cap AC_n(X) \text{ is homotopic to the identity mapping on } M_s \cap AC_n(X) \\ &\text{in } p_2^{-1}((0, 1)). \end{aligned}$$

We remark that by the flatness of H , if $H(M_s \cap AC_n(X), t) \subseteq p_2^{-1}((0, 1))$, then there is a neighborhood U of $(s; t)$ such that $H(M_{s'} \cap AC_n(X), t') \subseteq p_2^{-1}((0, 1))$ for any $(s'; t') \in U$. The following lemma can be proved similarly as [6, Lemma 4.3].

Lemma 4.2. *Let X be a noncontractible space and $H : AC_n(X) \times \mathbb{I} \rightarrow AC(X)$ a flat homotopy. If $H(M_0 \cap AC_n(X), t_0) \subseteq p_2^{-1}((0, 1))$, then there exists a neighborhood U of $(0; t_0)$ such that H does not satisfy $P(s, t)$ for any $(s; t) \in U$ with $s > 0$. An analogous statement holds for $H(M_1 \cap AC_n(X), t_0) \subseteq p_2^{-1}((0, 1))$.*

By Lemmas 4.1 and 4.2, we can use the proof of [6, Lemma 4.4] to verify also the next lemma.

Lemma 4.3. *Let X be a noncontractible space. If $H : AC_n(X) \times \mathbb{I} \rightarrow AC(X)$ is a flat homotopy such that $H(u, 0) = u$ for every $u \in AC_n(X)$, then $H(-, 1)$ is not a constant mapping.*

We can now regard the proof of [6, Theorem 1.2] also as a proof of Theorem 3.3. We shall only retrace its line of argument.

Proof of Theorem 3.3. Suppose that the space $AC(X)$ is contractible. Then there exists a homotopy $H : AC(X) \times T \rightarrow AC(X)$ connecting the identity mapping with the constant one. By compactness of the time interval \mathbb{I} , for every $a = (0; y) \in \{0\} \times \mathbb{I}$, there exists $\varepsilon_0 > 0$ such that the diameters of $p_2 \circ H(O_{\varepsilon_0}(a), t)$ are less than 1 for all $t \in \mathbb{I}$. Hence, by compactness of $[A, B]$, there exists $\varepsilon_1 > 0$ such that the diameters $p_2 \circ H(O_{\varepsilon_1}(a), t)$ are less than 1 for all $a = (0; y) \in \{0\} \times \mathbb{I}$ and all $t \in \mathbb{I}$. Let n be a number such that $1/n < \varepsilon_1$. By Lemma 4.1 we may then assume that $H|_{AC_n(X) \times \mathbb{I}}$ is a flat contraction. However, this contradicts Lemma 4.3. \square

5. PROOFS OF THEOREMS 3.4 AND 3.5

In this section we prove Theorems 3.4 and 3.5. Here we shall see a very different feature of $AC(X)$ in comparison with $SC(X)$.

Let $U = p_2^{-1}([0, 1))$ and $V = p_2^{-1}((0, 1])$. The spaces U, V and $U \cap V$ are homotopy equivalent to $p_2^{-1}(\{0\}), p_2^{-1}(\{1\})$ and $p_2^{-1}(\{1/2\})$, respectively. Let i_U and i_V be the inclusion maps from $U \cap V$ to U and V , respectively. Consider the following part of the Mayer-Vietoris exact sequence for the singular homology:

$$\begin{aligned} H_{n+1}(U) \oplus H_{n+1}(V) &\longrightarrow H_{n+1}(AC(X)) \longrightarrow H_n(U \cap V) \xrightarrow{(i_U^*, i_V^*)} \\ H_n(U) \oplus H_n(V) &\longrightarrow H_n(AC(X)). \end{aligned}$$

Since the inclusion maps from U to $AC(X)$ and from V to $AC(X)$ are inessential, we have the following exact sequence:

$$0 \longrightarrow H_{n+1}(AC(X)) \longrightarrow H_n(U \cap V) \xrightarrow{(i_U^*, i_V^*)} H_n(U) \oplus H_n(V) \longrightarrow 0.$$

We recall the following fact:

Lemma 5.1. [9, Theorem 1.3] *Let CX be the cone over a space X and $CX \vee CY$ the one-point union with two points of the base spaces X and Y being identified to a point. Then, the following holds for $n \geq 1$:*

$$H_n(X \vee Y) \cong H_n(X) \oplus H_n(Y) \oplus H_n(CX \vee CY).$$

Proof of Theorem 3.4. Let $r_X : X \vee Y \rightarrow X$ and $r_Y : X \vee Y \rightarrow Y$ be the retractions from Lemma 5.1. Then $H_n(CX \vee CY)$ is the kernel of the homomorphism $((r_X)_*, (r_Y)_*) : H_n(X \vee Y) \rightarrow H_n(X) \oplus H_n(Y)$.

$$\begin{array}{ccccc} V & \cong & p_2^{-1}(\{1\}) & = & p_2^{-1}(\{1\}) \\ \uparrow i_V & & & & \uparrow r_B \\ U \cap V & \cong & p_2^{-1}(\{1/2\}) & \cong & (p_2^{-1}(\{0\}), A) \vee (p_2^{-1}(\{1\}), B) \\ \downarrow i_U & & & & \downarrow r_A \\ U & \cong & p_2^{-1}(\{0\}) & = & p_2^{-1}(\{0\}) \end{array}$$

Since the diagram above is homotopy commutative, $H_{n+1}(AC(X))$ is isomorphic to $H_n(C(p_2^{-1}(\{0\})) \vee C(p_2^{-1}(\{1\})))$.

For each n , let X_n be a copy of the one-point union of X and the unit interval \mathbb{I} , where the attaching point is $0 \in \mathbb{I}$. Let x_n be a copy of $1 \in \mathbb{I}$. We regard X_n as $[C, C_n] \cup p^{-1}(\{C_n\})$. Then $p_2^{-1}(\{0\})$ is homotopy equivalent to $\bigvee_{n \in \mathbb{N}} (X_{2n-1}, x_{2n-1})$,

$p_2^{-1}(\{1\})$ is homotopy equivalent to $\tilde{V}_{n \in \mathbb{N}}(X_{2n}, x_{2n})$ and $p_2^{-1}(\{1/2\})$ is homotopy equivalent to $\tilde{V}_{n \in \mathbb{N}}(X_n, x_n)$. This verifies the assertion. \square

Proof of Theorem 3.5. By the result preceding the proof of Theorem 3.4, we have the exact sequence:

$$0 \longrightarrow H_2(AC(X)) \longrightarrow H_1(U \cap V) \xrightarrow{(i_{U*}, i_{V*})} H_1(U) \oplus H_1(V) \longrightarrow 0.$$

Suppose that $\pi_1(X)$ is trivial. Since $\pi_1(U \cap V)$ is isomorphic to $\prod_{n \in \mathbb{N}} \pi_1(X_n, x_n)$, $\pi_1(U \cap V)$ is also trivial. Hence $H_1(U \cap V)$ is trivial, which implies that $H_2(AC(X))$ is trivial, by the above exact sequence.

On the contrary, suppose that $\pi_1(X)$ is nontrivial. Choose a non-trivial element a of $\pi_1(X)$ and let $a_n \in \pi_1(X_n)$ be copies of a . Then there exists an element a^* of the group $\pi_1(U \cap V)$ such that its image under the induced homomorphism from retraction of the space $U \cap V$ to the $[C, C_{2n}] \cup p^{-1}([C_{2n}, C_1])$ is

$$[a_1, a_2][a_3, a_4] \cdots [a_{2n-1}, a_{2n}].$$

By [10, Theorem 1], the commutator length of this element is n . Since the number n is arbitrary, a^* does not belong to the commutator subgroup of $\pi_1(U \cap V)$. Therefore the homology class $[a^*] \in H_1(U \cap V)$ is nontrivial (2). However, its images $i_{U*}([a^*])$ and $i_{V*}([a^*])$ in $H_1(U)$ and $H_1(V)$, respectively, are obviously trivial. By the exactness of the Mayer-Vietoris sequence, the group $H_2(AC(X))$ is nontrivial. Since by Theorem 3.2, $\pi_1(AC(X))$ is trivial, it follows by the Hurewicz Theorem that $\pi_2(AC(X))$ is isomorphic to $H_2(AC(X))$ and hence is also nontrivial. \square

Proof of Theorem 3.6. Let \mathbb{H} be the well-known Hawaiian earring, which is obviously a Peano continuum. Let us show that $SC(\mathbb{H})$ is not homotopy equivalent to $AC(\mathbb{H})$. Consider the embedding $\varphi : \mathbb{H} \rightarrow SC(\mathbb{H})$, defined by $\varphi(x) = (A_1, x) \in SC(\mathbb{H})$ in Figure 1. In the sequel we shall identify \mathbb{H} with its image $\varphi(\mathbb{H})$.

We show that the mapping φ is not homotopy equivalent to the constant one. To show this by contradiction, suppose that this is not the case and that there exists a homotopy $H : X \times I \rightarrow SC(\mathbb{H})$ such that $H(x, 0) = \varphi(x)$ and $H(-, 1)$ is a constant mapping to some point $*$.

Since $SC(\mathbb{H})$ is a path connected space, we may assume without loss of generality, that $*$ $\in [A, B] \subset \mathcal{T}$. Let $a = \inf\{t : H(x^*, t) \in [A, B]\}$. The image $H(x^* \times [0, a])$ is a Peano continuum such that $H(x^*, 0) = (A_1, x^*) \in \mathcal{T} \subset SC(X)$, $H(x^*, 1) \in [A, B] \subset \mathcal{T} \subset SC(X)$. Since $[A, B]$ is a path connected component of the space \mathcal{T} , the set $p \circ H(x^* \times [0, a])$ is not a subset of \mathcal{T} and there exists $b \in [0, a]$ for which $p \circ H(x^*, b) \notin \mathcal{T}$ and consequently $H(x^*, b) \in \mathbb{I}^2 \setminus \mathcal{T}$.

Since $b < a$, we have $p \circ H(x^*, [0, b]) \cap [A, B] = \emptyset$ and so $H(x^*, [0, b]) \cap [A, B] = \emptyset$. There exists a neighborhood D of $H(x^*, b)$ which is a disk and is disjoint from \mathcal{T} . We have a neighborhood W of x^* for which $H(W \times [0, b]) \subset (SC(X) \setminus [A, B])$ and $H(W \times b) \subset D$. Since the disk D is contractible and the space \mathbb{H} is a retract of $SC(X) \setminus [A, B]$, we get a homotopy of W to \mathbb{H} connecting the embedding with a constant mapping. However, the embedding of W to \mathbb{H} is essential, which is a contradiction. Now we conclude that φ is an essential mapping.

Next we show that every mapping $\psi : \mathbb{H} \rightarrow AC(\mathbb{H})$ is inessential. First, observe that the embeddings of U and V to $AC(\mathbb{H})$ are inessential. For instance, for U , we have a deformation retraction of U to $p_y^{-1}(A)$ which can be contracted in $F(\mathbb{H})$ to a point. We have $\psi(x^*) \in U$ or $\psi(x^*) \in V$ and we only deal with the case $\psi(x^*) \in U$.

There exists a neighborhood \mathcal{O} of x^* such that $\psi(\mathcal{O}) \subset U$. All but finitely many circles of \mathbb{H} are contained in \mathcal{O} . Since $AC(X)$ is simply connected by Theorem 3.2, the restrictions of ψ to the finitely many circles are inessential mappings. Hence, as a total, ψ is an inessential map.

Suppose now that the spaces $SC(X)$ and $AC(X)$ are homotopy equivalent. Then there exist mappings $f : SC(X) \rightarrow AC(X)$ and $g : AC(X) \rightarrow SC(X)$ such that the composition gf is homotopic to the identity mapping. Thus $\varphi \simeq gf\varphi$. However, $gf\varphi$ is an inessential mapping, because $f\varphi$ is inessential and φ was shown to be an essential mapping. Contradiction. \square

In view of Theorems 3.2, 3.3, and 3.4, it is natural to ask the following question (cf. [7]):

Problem 5.2. *Does there exist a finite-dimensional noncontractible Peano continuum all homotopy groups of which are trivial?*

6. ACKNOWLEDGEMENTS

This research was supported by the Slovenian Research Agency grants P1-0292-0101, J1-9643-0101, J1-2057-0101 and J1-4144-0101. The first author was also supported by the Grant-in-Aid for Scientific Research (C) of Japan No. 20540097 and 23540110. We thank the referee for comments and suggestions.

REFERENCES

- [1] M. G. Barrat, J. Milnor, *An example of anomalous singular homology*, Proc. Amer. Math. Soc. **13** (1962), 293–297.
- [2] K. Borsuk, *Theory of Shape*, Monografie Math. **59**, PWN, Warsaw, 1975.
- [3] G. E. Bredon, *Sheaf Theory*, Second Ed., Graduate Texts in Math. **170**, Springer, New York, 1997.
- [4] K. Eda, *Free σ -products and noncommutatively slender groups*, J. Algebra **148** (1992), 243–263.
- [5] K. Eda, U. H. Karimov, D. Repovš, *On (co)homology locally connected spaces*, Topology Appl. **120** (2002), 397–401.
- [6] K. Eda, U. H. Karimov, D. Repovš, *A construction of simply connected noncontractible cell-like two-dimensional continua*, Fund. Math. **195** (2007), 193–203.
- [7] K. Eda, U. H. Karimov, D. Repovš, *A nonaspherical cell-like 2-dimensional simply connected continuum and related constructions*, Topology Appl. **156** (2009), 515–521.
- [8] K. Eda, U. H. Karimov, D. Repovš, *On the second homotopy group of $SC(Z)$* , Glasnik Math. **44**(64):2 (2009), 493–498.
- [9] K. Eda and K. Kawamura, *Homotopy groups and homology groups of the n -dimensional Hawaiian earring*, Fund. Math. **165** (2000), 17–28.
- [10] H. B. Griffiths, *A note on commutators in free products. II*, Proc. Cambridge Philos. Soc. **51** (1955), 245–251.
- [11] U. H. Karimov, D. Repovš, *On suspensions of contractible compacta of trivial shape*, Proc. Amer. Math. Soc. **127**:2 (1999), 627–632.
- [12] R. C. Lyndon, P. E. Schupp, *Combinatorial Group Theory*, Princeton Univ. Press, Princeton, N.J., 1971.
- [13] E. H. Spanier, *Algebraic Topology*, Springer-Verlag, New York, Heidelberg, Berlin, 1966.
- [14] G. W. Whitehead, *Elements of Homotopy Theory*, Springer-Verlag, Berlin, 1978.

SCHOOL OF SCIENCE AND ENGINEERING, WASEDA UNIVERSITY, TOKYO 169-8555, JAPAN

E-mail address: `eda@logic.info.waseda.ac.jp`

INSTITUTE OF MATHEMATICS, ACADEMY OF SCIENCES OF TAJIKISTAN, UL. AINY 299^A, DUSHANBE
734063, TAJIKISTAN

E-mail address: `umedkarimov@gmail.com`

FACULTY OF MATHEMATICS AND PHYSICS, AND FACULTY OF EDUCATION, UNIVERSITY OF LJUBLJANA,
P.O.BOX 2964, LJUBLJANA 1001, SLOVENIA

E-mail address: `dusan.repovs@guest.arnes.si`